

CRESTON IPMA INVASIVE PLANT (TERRESTRIAL AND RIPARIAN) PRIORITY LIST– 2017

(* not known in CKISS region; BC=biocontrol)

Regional EDRR - Not currently known in this IPMA (and may also not be known in the CKISS region*). Follow EDRR reporting and action protocols. Bolded species are EDRR for the province.		
<ul style="list-style-type: none"> - Bighead knapweed - Black knapweed - Buffalobur* - Caraway - Coltsfoot - Field scabious - Flowering rush* - Garlic mustard* 	<ul style="list-style-type: none"> - Giant hogweed - Gorse* - Greater celandine - Greater knapweed - Hoary cress - Longspine sandbur* - Marsh plume thistle 	<ul style="list-style-type: none"> - Nodding thistle* - Puncturevine* - Russian knapweed* - Tansy ragwort - Wild chervil - Wild four o'clock* - Wood sage*
ERADICATION or ANNUAL CONTROL – Species known in IPMA but with very limited distribution. Enter inventory data, report and treat or monitor annually.		
<ul style="list-style-type: none"> - ALL containment species OUTSIDE containment lines - Baby's breath - Black locust - Common bugloss 	<ul style="list-style-type: none"> - Leafy spurge - Meadow/brown/black knapweed - Plumeless thistle - Policeman's helmet 	<ul style="list-style-type: none"> - Purple loosestrife - Rush skeletonweed - Scotch thistle - Teasel
CONTAINMENT – Enter inventory data, report and treat all sites outside containment lines. Some of these species have biocontrol available which can be used within the containment line (BC).		
<ul style="list-style-type: none"> - Blueweed (contain to Wynndel) - Common tansy (keep out of E. Kootenay – Crackerjack FSR east) - Himalayan blackberry (prevent from spreading east of Kitchener) - Hoary alyssum (contain to W. of Erickson) 	<ul style="list-style-type: none"> - Knotweeds (contain to sites that do not border natural lands, transportation corridors, or riparian areas) - Scotch broom (contain to Kuskanook north) - Spotted knapweed (contain to west of Yahk gas line) - Yellow flag iris (contain to CVWMA and gardens) 	Contain to gardens: <ul style="list-style-type: none"> - Butterfly bush - Cypress spurge - English ivy - Field bindweed - Goutweed - Japanese butterbur - Lamium/Yellow archangel - Russian olive - Periwinkle - Salt cedar - Siberian elm
ESTABLISHED (BIOCONTROL OR SITE-SPECIFIC APPROACH) – Widespread species that are beyond landscape-level control or have relatively low impact. Treat based on specific land management objectives.		
<ul style="list-style-type: none"> - Bull thistle (BC) - Burdock - Canada thistle (BC) - Chicory - Curled dock - Dalmatian toadflax (BC) 	<ul style="list-style-type: none"> - Dame's rocket - Diffuse knapweed - Hairy cat's ear - Hound's tongue (BC) - Mullein - Orange hawkweed 	<ul style="list-style-type: none"> - Oxeye daisy - Scentless chamomile - St. John's wort (BC) - Sulphur cinquefoil - Yellow hawkweeds - Yellow/common toadflax
WATCHLIST – There is a lack of information on the distribution and density of this species within this IPMA.		
<ul style="list-style-type: none"> Sea buckthorn Queen Anne's Lace Wormwood (Absinth) Yellow bedstraw 		

