

LOWER ARROW PEND D'OREILLE IPMA INVASIVE PLANT PRIORITY LIST – 2017

(* not known in CKISS region; BC=biocontrol; **species distribution/density to be verified in 2017)

Regional EDRR - Not currently known in this IPMA (and may also not be known in the CKISS region*). Follow EDRR reporting and action protocols.		
<ul style="list-style-type: none"> - Bighead knapweed - Buffalobur* - Caraway - Coltsfoot* - Common bugloss - Cypress spurge* - Flowering rush* - Garlic mustard* 	<ul style="list-style-type: none"> - Giant hogweed - Gorse* - Greater celandine - Leafy spurge (one site in IPMA but managed by FLNRO) - Longspine sandbur* - Marsh plume thistle - Meadow/brown/black knapweed 	<ul style="list-style-type: none"> - Nodding thistle* - Puncturevine* - Russian knapweed* - Scentless chamomile - Tansy ragwort - Wild four o'clock* - Wood sage*
ERADICATION or ANNUAL CONTROL – Species known in IPMA but with very limited distribution. Enter inventory data, report and treat or monitor annually.		
<ul style="list-style-type: none"> - Containment species OUTSIDE lines and: - Baby's breath - Blueweed 	<ul style="list-style-type: none"> - Greater knapweed - Purple loosestrife - Rush skeletonweed - Scentless chamomile 	<ul style="list-style-type: none"> - Scotch broom - Scotch thistle - Teasel - Wild chervil**
CONTAINMENT – Enter inventory data, report and treat all sites outside containment lines. Some of these species have biocontrol available which can be used within the containment line (BC).		
<ul style="list-style-type: none"> - Field scabious (contain to Salmo) - Himalayan blackberry (prevent from spreading north of Cayuse Cr.) - Hoary alyssum (prevent from spreading north of Deer Park) 	<ul style="list-style-type: none"> - Hoary cress (contain to Warfield and Waneta/Nelway) - Policeman's helmet (contain to Trail/Rossland) - Yellow flag iris (contain to Meldeanna pond) 	<p>Contain to gardens:</p> <ul style="list-style-type: none"> - Butterfly bush - Cypress spurge - English ivy - Field bindweed - Goutweed - Japanese butterbur - Lamium/Yellow archangel - Periwinkle - Russian olive - Salt cedar - Siberian elm
ESTABLISHED (BIOCONTROL OR SITE-SPECIFIC APPROACH) – Widespread species that are beyond landscape-level control or have relatively low impact. Treat based on specific land management objectives.		
<ul style="list-style-type: none"> - Black locust - Bull thistle (BC) - Burdock - Canada thistle (BC) - Chicory - Common tansy - Curled dock - Dalmatian toadflax (BC) 	<ul style="list-style-type: none"> - Dame's rocket - Diffuse knapweed (BC) - Hairy cat's ear - Hound's tongue (BC) - Mullein - Orange hawkweed - Oxeye daisy - Plumeless thistle (BC) 	<ul style="list-style-type: none"> - Spotted knapweed (BC) - St. John's Wort (BC) - Sulphur cinquefoil - Queen Anne's lace - Yellow/common toadflax - Yellow hawkweeds
WATCHLIST – There is a lack of information on the distribution and density of this species within this IPMA.		
<ul style="list-style-type: none"> Wormwood Yellow bedstraw 		