

KASLO NORTH IPMA INVASIVE PLANT (TERRESTRIAL AND RIPARIAN) PRIORITY LIST – 2017

(* not known in CKISS region; BC=biocontrol)

Regional EDRR - Not currently known in this IPMA (and may also not be known in the CKISS region*). Follow EDRR reporting and action protocols. Bolded species are EDRR for the province.		
<ul style="list-style-type: none"> - Baby's breath - Bighead knapweed - Buffalobur* - Caraway* - Common bugloss* - Field scabious - Flowering rush* - Garlic mustard* - Gorse* 	<ul style="list-style-type: none"> - Greater celandine - Greater knapweed - Hoary cress - Leafy spurge - Longspine sandbur* - Marsh plume thistle - Meadow/brown/black knapweed - Nodding thistle* - Plumeless thistle 	<ul style="list-style-type: none"> - Puncturevine* - Rush skeletonweed - Russian knapweed* - Scentless chamomile - Scotch thistle - Teasel - Wild four o'clock* - Wood sage* - Yellow flag iris - Tansy Ragwort
ERADICATION or ANNUAL CONTROL – Species known in IPMA but with very limited distribution. Enter inventory data, report and treat or monitor annually. Some of these species have biocontrol (BC) available.		
<ul style="list-style-type: none"> - ALL containment species OUTSIDE their containment lines (see next section) 	<ul style="list-style-type: none"> - Blueweed - Giant hogweed - Hoary alyssum 	<ul style="list-style-type: none"> - Policeman's helmet - Purple loosestrife
CONTAINMENT – Enter inventory data, report and treat all sites outside containment lines. Some of these species have biocontrol (BC) available which can be used within the containment line.		
<ul style="list-style-type: none"> - Coltsfoot (contain to Kaslo) - Himalayan blackberry (prevent from spreading north of Kaslo) - Knotweeds (contain to sites that do not border natural lands, transportation corridors, or riparian areas) 	<ul style="list-style-type: none"> - Scotch broom (prevent from spreading north of Shutty Bench) <p>Contain to gardens:</p> <ul style="list-style-type: none"> - Butterfly bush - Cypress spurge - English ivy - Field bindweed 	<p>Contain to gardens:</p> <ul style="list-style-type: none"> - Goutweed - Japanese butterbur - Lamium/Yellow archangel - Russian olive - Periwinkle - Salt cedar - Siberian elm
ESTABLISHED (BIOCONTROL OR SITE-SPECIFIC APPROACH) – Widespread species that are beyond landscape-level control or have relatively low impact. May have biocontrol (BC) available. Treat based on land management objectives.		
<ul style="list-style-type: none"> - Bull thistle (BC) - Burdock - Canada thistle (BC) - Chicory - Common tansy - Curled dock - Dalmatian toadflax (BC) 	<ul style="list-style-type: none"> - Dame's rocket - Diffuse knapweed (BC) - Hairy cat's ear - Hound's tongue (BC) - Mullein - Orange hawkweed - Oxeye daisy 	<ul style="list-style-type: none"> - Spotted knapweed (BC) - St. John's wort (BC) - Sulphur cinquefoil - Yellow hawkweeds - Yellow/common toadflax
WATCHLIST – There is a lack of information on the distribution and density of this species within this IPMA.		
<ul style="list-style-type: none"> Black locust Bur chervil Comfrey Meadow salsify Wall lettuce 	<ul style="list-style-type: none"> Wild carrot/Q. Anne's Lace Wild chervil Wormwood (Absinth) Yellow bedstraw 	

