

NELSON IPMA INVASIVE PLANT (TERRESTRIAL AND RIPARIAN) PRIORITY LIST – 2017

(* not known in CKISS region; BC=biocontrol)

Regional EDRR - Not currently known in this IPMA (and may also not be known in the CKISS region*). Follow EDRR reporting and action protocols. Bolded species are EDRR for the province.		
<ul style="list-style-type: none"> - Bighead knapweed - Black knapweed* - Buffalobur* - Caraway* - Coltsfoot - Common bugloss* - Flowering rush* - Garlic mustard* - Giant hogweed 	<ul style="list-style-type: none"> - Gorse* - Greater celandine - Greater knapweed - Hoary cress - Leafy spurge - Longspine sandbur* - Marsh plume thistle* - Meadow/brown knapweed - Nodding thistle* 	<ul style="list-style-type: none"> - Plumeless thistle - Puncturevine* - Russian knapweed* - Tansy ragwort - Wild chervil - Wild four o'clock* - Wood sage*
ERADICATION or ANNUAL CONTROL – Species known in IPMA but with very limited distribution. Enter inventory data, report and treat or monitor annually. Some of these species have biocontrol (BC) available.		
<ul style="list-style-type: none"> - ALL containment species OUTSIDE containment lines) - Baby's breath - Blueweed 	<ul style="list-style-type: none"> - Black knapweed - English ivy - Mouse-ear hawkweed 	<ul style="list-style-type: none"> - Rush skeletonweed - Scentless chamomile - Scotch thistle - Teasel
CONTAINMENT – Enter inventory data, report and treat all sites outside containment lines. Some of these species have biocontrol (BC) available which can be used within the containment line.		
<ul style="list-style-type: none"> - Field scabious (contain to Salmo) - Knotweeds (contain to sites that do not border natural lands, transportation corridors, or riparian areas) - Policeman's helmet (contain to Blewett) - Purple loosestrife (contain 4-Mile to Kootenay Canal) 	<ul style="list-style-type: none"> - Scotch broom (contain from spreading south and west of Nelson) - Yellow flag iris (contain to 4-mile to Bird Creek Marsh) <p>Contain to gardens:</p> <ul style="list-style-type: none"> - Butterfly bush - Cypress spurge - English ivy - Field bindweed 	<p>Contain to gardens:</p> <ul style="list-style-type: none"> - Goutweed - Japanese butterbur - Lamium/Yellow archangel - Russian olive - Periwinkle - Salt cedar - Siberian elm
ESTABLISHED (BIOCONTROL OR SITE-SPECIFIC APPROACH) – Widespread species that are beyond landscape-level control or have relatively low impact. May have biocontrol (BC) available. Treat based on land management objectives.		
<ul style="list-style-type: none"> - Bladder campion - Bull thistle (BC) - Burdock - Canada thistle (BC) - Chicory - Common tansy - Curled dock - Flat peavine 	<ul style="list-style-type: none"> - Dalmatian toadflax (BC) - Dame's rocket - Diffuse knapweed (BC) - Hairy cat's ear - Himalayan blackberry - Hoary alyssum - Hound's tongue - Mullein 	<ul style="list-style-type: none"> - Orange hawkweed - Oxeye daisy - Spotted knapweed (BC) - St. John's wort (BC) - Sulphur cinquefoil - Yellow hawkweeds - Yellow/common toadflax
WATCHLIST – There is a lack of information on the distribution and density of this species within this IPMA.		
<ul style="list-style-type: none"> Queen Anne's lace Black locust 	<ul style="list-style-type: none"> Bristly locust 	